

# Stress-induced changes of cytoplasmic free calcium in lectin-simulated mouse T lymphocytes

P. CSERMELY, S. TÓTH \*, E. BEREGI \*  
and M. BOROSS \*

*Institute of Biochemistry I  
\* Gerontology Center  
Semmelweis University Budapest (H)*

The XVth Congress  
of the International  
Association of  
Gerontology

Budapest, Hungary  
4-9 July 1993

## SUMMARY

Extreme environmental or homeostatic changes provoke an adaptive stress-response of the organism. In our present work we have investigated the effect of "overcrowd-stress" on the Concanavalin-A stimulated cytoplasmic free Ca concentration of mouse splenic T lymphocytes. Mice were kept under "normal" (68 cm<sup>2</sup>/animal) or "overcrowded" (22 cm<sup>2</sup>/animal) conditions for 10 days. No change was observed in resting Ca after overcrowd-stress. The lectin-induced rise in intracellular Ca level of splenic T lymphocytes, however, was three times higher ( $p < 0.01$ ) in stressed mice compared to the control group.

## INTRODUCTION

A significant rise in intracellular calcium concentration is one of the first signals of T lymphocyte activation [1,2]. Overcrowding and other forms of psychosocial stress were shown to modulate the immune response in mice and rats [3-6]. In our present study we have investigated the effect of "overcrowd-stress" on resting and stimulated cytoplasmic free Ca concentration of mouse splenic T lymphocytes.

## MATERIALS AND METHODS

### *Reagents and cells*

Concanavalin-A (type IV), digitonin, dimethyl-sulfoxide (DMSO), EGTA,