

Chapter XVII

EXPECTED FUNCTIONS OF SALIVARY HSP70 IN THE ORAL CAVITY (REVIEW ARTICLE)

***Tibor Károly Fábián¹, Csaba Sőtér², Minh Tu Nguyen², Péter
Csermely² and Pál Fejérdy¹***

¹Semmelweis University Budapest, Faculty of Dentistry, Clinic of Prosthetic Dentistry,
Budapest, Hungary, EU;

²Semmelweis University Budapest, Faculty of Medicine, Institute of Medical Chemistry
Molecular Biology and Pathobiochemistry, Budapest, Hungary, EU.

ABSTRACT

Saliva is a major determinant of the environment on all oral surfaces in both healthy and pathological conditions. It is a body fluid, secreted by three pairs of major salivary glands and by many of minor salivary glands supplemented with several constituents originated from intact or destroyed mucosal cells, immune cells, oral microorganism and blood. *Salivary Hsp70* is an important constituent of saliva, originating from several sources including salivary glands, mucosal cells, gingival crevicular fluid, blood and oral microbes. Salivary Hsp70 seems to be excreted constitutively, but its level can be increased significantly with psychological stress conditions, and also with chewing and taste stimulation. Several physiotherapeutic procedures like massage and local heat stimulation of major salivary glands also increases the amount. *Expected functions* of salivary Hsp70 include immunological and other antimicrobial defense mechanisms, and also cytoprotective effects on mucosal and immune cells. Consequently, it likely plays a role in healing of mucosal wounds and ulcers; in defense against gingival inflammation; in release of immune answer toward mucosal tumors; and also in allergic and autoimmune mucosal processes. Salivary Hsp70 is also likely participating in the formation of acquired pellicle on tooth and denture surfaces, leading to a possible role in physico-chemical defense of tooth surfaces; in crystal growth homeostasis of the teeth; in bacterial adhesion to tooth/denture surfaces; and in prevention and healing of denture

induced mucosal irritations. However our knowledge related to salivary Hsp70 is still rather limited. Consequently the expectations about several functions detailed in this chapter are partially based on data related to extracellular but not particularly salivary Hsp70.

INTRODUCTION

Saliva and the Oral Cavity

Saliva is a body fluid, secreted by three pairs of major salivary glands (parotid, submandibular and sublingual) and by many of minor salivary glands [28]. Primary saliva is secreted in secretory endpieces (acini) of salivary glands. Primary saliva is modified in the intercalated, striated and excretory (collecting) ducts leading from the acini to the mouth [116]. In the intercalated duct water and electrolyte transport into the saliva is supposed. Striated duct is responsible for both electrolyte secretion and reabsorption. Probably there is a transport of proteins like IgA, lysozyme, kallikrein and Hsp70 [28,29] in the striated duct as well. Some electrolyte transport in the excretory (collecting) duct is also suspected [116]. Primary saliva is also modified by serum exudates via tight junctions between several glandular cells (ultrafiltration) and via transcellular diffusion through these cells [37,51]. Entering the mouth, ductal saliva of several salivary glands are blended, and supplemented with many constituents originated from intact or destroyed mucosal cells, immune cells, and oral microorganism. Blood constituents also enter the oral cavity via gingival crevicular fluid, via the mucosa as mucosal transudate, and via intraoral bleeding [28]. Consequently a complex mixture of a high variety of molecules is resulted in the oral cavity, frequently called as "mixed saliva" and/or "whole saliva" in the scientific literature [28].

Main Functions of Saliva

Saliva is a major determinant of environment on all the oral surfaces. On tooth surfaces saliva plays an important role in acquired pellicle formation, which is a thin (ca. 0.5-1 μ m) layer of several salivary proteins with calcium hydroxide binding properties [15,48]. Acquired pellicle plays a major role in crystal growth homeostasis of the teeth, and in physico-chemical defense of tooth surfaces. Beside these, acquired pellicle plays a major role in bacterial adhesion on the tooth surfaces [15,48]. Saliva also strongly influences the bacterial biofilm formation following adhesion of bacteria on pellicle [28]. Similarly, saliva plays an important role in physico-chemical defense of the oral (and upper gastro intestinal) soft tissues. Its role is also crucial in antimicrobial defense mechanisms, and in wound (and ulcer) healing pathways of the oral cavity [28]. Saliva is also highly important for appropriate taste-sensation [58].

The Hsp70 Chaperone Family

The human heat shock protein 70 (Hsp70) chaperone family contains at least eight homologous chaperone proteins. Endoplasmic reticulum (ER) and mitochondria have their specific Hsp70 proteins. The remaining six family members reside mainly in the cytosol and nucleus, but some of them are also present in lysosomes [26]. There are also several homologues in bacteria (i.e. DnaK in *E. coli*) and fungi (i.e. Ssa1p/Ssa2p in *C. albicans*). Although there are some significant differences, all known members of human Hsp70 family display highly conserved amino acid sequences, and domain structures. All these proteins consist of an ATPase domain at the N-terminal, followed by a protease sensitive region, a peptide binding domain, and a variable region [26]. Hsp70 molecules present in the cytosol, nucleus and lysosomes also contain a C-terminal region with a certain amino acid motif enabling the proteins to bind co-chaperones and other Hsps. The amino acid sequences of Hsp70 molecules localized in mitochondria and endoplasmic reticulum start with a unique N-terminal localization signal for delivery to the right cellular compartment. Hsp70 molecules in the endoplasmic reticulum have also a C-terminal retention signal sequence, which inhibits their exit from the lumen [26]. Each members of Hsp70 family and their most important intracellular functions are briefly summarized as follows.

Hsp70-1 (*Hsp70*, *Hsp72*) is stress inducible protein with two isoforms (*Hsp70-1a,b*). It can be found in the cytosol, in the nucleus, and also on the luminal side of the lysosomal membrane. Its main function is to cope with harmful aggregations of denatured proteins via chaperoning function [34,45,70], and stabilize lysosomal membranes during stress [26,44]. *Hsp70-1* was also showed to influence signal transduction pathways, cell cycle and senescence regulation [80,94,97]. *Hsp70-1t* (*Hsp70-hom*) is a constitutively expressed protein, localized in the cytosol and nucleus. It is expressed mainly in testis. Its specific role is not yet clear [26]. *Hsp70-2* (*Hsp70-3*, *HspA2*) is also constitutively expressed protein localized in the cytosol and nucleus. It is expressed mainly in testis and brain. It is involved in senescence and cell cycle regulation and inhibition of apoptotic pathways [26,94,97]. *Hsp70-5* (*Bip*; *Grp78*) is a constitutively expressed and roughly endoplasmic reticulum specific protein. Its main function is to facilitate the transport of newly synthesized proteins into the ER lumen and their subsequent folding [110]. It also alleviates ER stress induced apoptotic stimuli via Ca^{2+} binding and preventing Ca^{2+} efflux from the ER [39,79]. *Hsp70-6* (*Hsp70B'*) is a stress inducible protein which is localized in the cytosol and nucleus with similar functions like *Hsp70-1*. It is expressed mainly in dendritic cells, monocytes and NK cells [26]. *Hsp70-8* (*Hsc70*; *Hsp73*, *Hsp70-cognate*) is a constitutively expressed protein in most tissues. It is involved in multiple housekeeping chaperoning functions including protein folding and refolding, protein translocation across membranes, chaperone-mediated autophagy, prevention of protein aggregation and disassemble of clathrin coated vesicles [26,34,45]. *Hsp70-9* (*mtHsp70*, *mortalin*, *Grp75*) is a constitutively expressed and roughly mitochondria specific protein. Its main function is to assist the incoming proteins in correct folding following transmembrane transport, but outside mitochondria it can also regulate senescence pathways [27,97].

Main Extracellular Functions of Hsp70

In addition to intracellular response, stress also triggers release of Hsp70 into the extracellular space, and Hsp70 is also released from cells undergoing necrosis [17]. Extracellular Hsp70 reveal cytoprotective properties through cell surface association [49], which may be followed by internalization [42,43]. Extracellular Hsp70 is also involved in a number of physiological and pathological events including modulation of cytokine release [2,3], immunity [68,71,89,90,100], and the modulation of neuronal function [16,17,21,43,106]. In addition, Hsp70 is able to enter the blood stream [86,93,117], and possess the ability to act at distant sites of the body as an ancestral danger signal [65,66] triggered by cell injury, immune-inflammatory reactions, and physical [32,57] or behavioral [31,35] stress of the organism. The presence of Hsp70 in human saliva was also demonstrated [30,31] indicating that, premised extracellular functions of Hsp70 may play an important role on the mucosal, periodontal and tooth surfaces too, as detailed in the next sections of this chapter.

ORIGIN AND SUBTYPES OF SALIVARY HSP70

Salivary Gland Origin

Salivary glands are one of the main sources of Hsp70 in the saliva [29,30,31]. Although it has not yet been investigated in detail, it is very likely that, Hsp70 of glandular origin is not a new member of Hsp70 family, but rather a mixture of several known Hsp70-type proteins including stress inducible Hsp70-1 (Hsp72) and constitutive Hsp70-8 (Hsp73) [29,30,31]. The level of salivary Hsp70 shows rather big differences between subjects in both mixed (whole) saliva and cannulated parotid duct saliva (authors' unpublished data). There are also within subject differences may be because salivary Hsp70 is prompt inducible via several stimuli (see also below) [30,31,33]. Hsp70 is not secreted due to the classical secretory exocytotic process of the acinar cells, since the secretory parameters of Hsp70 were independent of those of amylase (a known secretory protein of salivary gland acinar cells) [30,31]. The transport of Hsp70 may involve passive transport via the salivary glands from blood serum [29,30,31], similarly to other blood proteins excreted with saliva [28,37,51]. A small capacity active transport from the striated duct cells of human salivary glands is also possible [29,30,31], since a higher amount of Hsp70 was detected in these cells comparing to other cell types of salivary glands and ducts [109]. Such transport would require either increased cell membrane permeability, or a specific exocytotic pathway [32,57], because Hsp70 family lacks a conventional signal peptide sequence that would allow them to be transported through cytosolic or plasma membranes [41]. An alternative transport of Hsp70 may occur either through lipid rafts [14,20,56], or through exosomes [22,55,56]. Another alternative transport mechanism was reported through lysosomes releasing their content into the extracellular space [17,63].

Mucosal Cell Origin

Mucosal cells are other main source of salivary Hsp70 [29]. Depending on location the oral epithelium can be either keratinized or non-keratinized. Keratinized surfaces (i.e. attached gingiva, hard palate, most dorsal surfaces of tongue) are subjected to strong mechanical loads. The abraded superficial cells of these surfaces lack intracytoplasmic organelles, being filled with keratin. In contrast, mucosal cells with more or less intact cell organelles are continuously released from the surface of moderately stressed non-keratinized epithelium (i.e. labial-, buccal-, alveolar mucosa, floor of the mouth, ventral surface of tongue) [9]. Hence a mixture containing intact and/or fragmented cells, intracellular compartments, membrane particles, and usual cytosolic constituents are released into the saliva from the non-keratinized mucosal surfaces [28]. Consequently all known members of Hsp70 family regularly present in human cells are released into the saliva from this mucosal source.

Gingival Crevicular Fluid Origin

Gingival crevicular fluid is also a main source of salivary Hsp70 [29]. From this source serum exudate and some intact and/or destroyed immune cells can reach saliva, also in healthy and particularly in inflamed gingival (and/or periodontal) conditions [28,51]. Consequently Hsp70 family members present in the blood serum and immune cells enters saliva from this source.

Blood Origin

Beside serum exudates originated from the gingival crevicular fluid and salivary glands (see above), serum exudate also reaches saliva through the oral mucosa as mucosal transudate [28,51]. Moreover, each of the cellular and other blood constituents may enter saliva via intraoral bleeding (i.e. bleeding periodontal pockets, wounds, ulcers) [28,51]. Hsp70 family members present in blood serum and blood cells reaches saliva from these sources.

Bacterial (Microbial) Origin

Other main sources of salivary Hsp70 are oral bacteria and other oral microbes [29]. But it should be noted that, bacterial (microbial) homologues of Hsp70 family are originated from this sources only [29]. The amount of microbial Hsp70 homologues in the saliva may be dependent on oral hygiene conditions, properties of bacterial biofilm on tooth surfaces, and microbial flora of the oral cavity and periodontal pockets (if any).

SALIVARY HSP70 AND SURFACE DEFENSE

Defense of Mucosal Surfaces

There are two major facets of mucosal defense mechanisms expected for salivary Hsp70. One of them involves immunological and other antimicrobial defense pathways [29] and will be detailed later (see below). The other is based on the cytoprotective effects of extracellular Hsp70 [42,43,49]. The cytoprotective effects seem to be based on three different mechanisms, such as aspecific binding of Hsp70 on mucosal cell surfaces [49]; a more specific adhesin-type binding to sulfoglycolipid structures of mucosal cells [12]; and surface receptor binding of Hsp70 mostly followed by internalization [17,42,43]. Aspecific binding leads to surface defense against toxins [49] very likely through protection and repair (chaperoning) of mucosal cell surface proteins [30,31]. Adhesin-type binding may prevent bacterial colonization of mucosal surfaces through occupying mucosal binding sites of Hsp70 related bacterial adhesins [12,92]. Surface receptor binding of Hsp70 was proved to several receptors like oxidized low density lipoprotein binding receptor CD91 [5]; tumor necrosis factor receptor CD40 [8]; toll like pattern recognition receptors TLR2 and TLR4 [3]; chemokine receptor CCR5 [36]; and scavenger receptors LOX-1, SREC-1 and FEEL-1/CLEVER-1 [102]. Although premised Hsp70 receptor bindings were studied mostly related to immunological context, it should be considered that, other cell types are also capable of receptor mediated binding and uptake of Hsp70 [17,21,43]. Up to the present, decrease of the cells' apoptotic and necrotic liability [42] and release of several cytokines [2,3] are the most important proved mechanisms of Hsp70 receptor induced cytoprotection. However, cytoprotective mechanism might be varied, since Hsp70 likely recognize different receptors on different cell types [103], and different members of Hsp70 family releases different effects on the same receptor [17]. The cellular answer may also be influenced by interactions of several Hsp70 receptors of the same cell surface [17] resulting an even more complex picture of possible cytoprotective pathways.

Defense of Tooth Surfaces

Recent data indicated that, Hsp70-1 (inducible Hsp72) binds both Gram-positive (*Streptococcus mutans* and *mitis*) and Gram-negative (*Escherichia coli*) bacteria [78]. Similarly, salivary Hsp70 also binds *S. mutans*, *S. mitis* and *E. coli*, moreover binds hydroxyl apatite the major inorganic component of tooth surfaces, (authors unpublished data). Taking together these results, it is likely that, salivary Hsp70 may play a role in the acquired pellicle formation followed by bacterial adhesion on tooth surfaces. The capability to take part in the acquired pellicle formation and to bind bacteria refers to that; salivary Hsp70 in general facilitates bacterial colonization of tooth surfaces. But it should be considered that, the pH optimum of bacterial binding of salivary Hsp70 is rather acidic (in our unpublished data it is around pH 5.0 - 5.5 similarly to recently published data with Hsp70-1 [78]). Such a low pH is not typical for tooth surfaces at the early stage of bacterial colonization, consequently salivary Hsp70 rather inhibits than facilitates bacterial adhesion to the acquired pellicle.

Recent finding also indicated that, ATP partially inhibits bacterial binding of Hsp70-1 [78]. Similarly, binding of salivary Hsp70 to bacteria and also to hydroxyl apatite was found to be partially inhibited by ATP (authors' unpublished data). The meaning of these latter ATP connected data is not yet clear but should be considered in future research of acquired pellicle formation and bacterial adhesion.

Immunological and Antimicrobial Defense Mechanisms

Three major facets of immune activation have been described for Hsp70 [69]. *The first of these* involves the appearance of intracellular Hsp70 on the surface of certain tumor cells leading to lysis of these cells by natural killer (NK) cells [11,72,73]. *The second facet* of immune activation involves released extracellular Hsp70 as an ancestral danger signal [65, 66] of cellular stress, death or lysis. Importantly both, uncomplexed ("free") and membrane bound (lipid rafts, exosomes) Hsp70 was showed to express such danger signal properties [73]. The immune activation here is very similar to that of bacterial lipopolysaccharides (LPS) and the effect of LPS and extracellular Hsp70 seems to be additive [19]. Hsp70 as a danger signal induces: release of proinflammatory cytokines from several immune cells (i.e. monocytes, dendritic cells, macrophages, T lymphocytes) [2,13,19,115]; release of NO from macrophages [83]; activation of NK cells [68,71]; activation of complement via an antibody-independent alternative pathway [90]. *The third facet* of immune system activation involves complexes of extracellular Hsp70 and other peptides. Because its chaperoning ability, uncomplexed Hsp70 binds other peptides, and as complex induces receptor-mediated uptake into antigen-presenting cells (i.e. macrophages, Langerhans and dendritic cells) to cross present this complex as an antigen (coupled with MHC-I or MHC-II molecules) to cytotoxic T cells and NK cells [10,101,107]. This mechanism is important in the defense against bacteria (and other microbes) and also as an initiator of immune defense against tumor cells and virus infected cells. From premised three major immunological facets, only the first one is not expected for salivary Hsp70 because in this case the tumor cells express their own Hsp70 on the surface. The other two functions are expected for salivary Hsp70 [29], since oral mucosa is extensively populated by antigen presenting Langerhans and dendritic cells [25]. (The expectation is also confirmed by the interesting finding that, Langerhans cells are properly oriented to "sample" the oral fluids with their dendrites toward the surface [25,47].)

Beside the above most intensely researched immunological functions, there is another immunological mechanism, in which extracellular Hsp70 takes part. Recent data indicate that, Hsp70-1 (inducible Hsp72) exert an opsonizing effect on bacteria, which activate the killing activity of polymorphonuclear neutrophil (PMN) leukocytes [78]. Although there is a considerable flux of neutrophils through the gingival sulcus into the saliva even in health, this function may be especially effective under inflammatory conditions (gingival inflammations, wound healing, fever), because Hsp70 binding of bacteria is facilitated by acidic milieu (pH 5.5) and high temperature (42°C) [78]. It should be noted that, opsonizing effect similar to that of Hsp70-1 [78] is not yet verified with salivary Hsp70. However it is likely because the bacterial binding property of salivary Hsp70 is very similar to that of Hsp70-1: it is also

facilitated by acidic milieu (pH 5.0-5.5) and high (42°C) temperature (authors' unpublished data, see also above).

Beside immunological pathways there are two other hypothesized antimicrobial defense mechanisms of salivary Hsp70 which should be considered. One of them is based on the bacterial binding property of Hsp70 mentioned above. Via this binding salivary Hsp70 may also entrap and agglutinate bacteria. Since Hsp70 is known to be able to form dimers and oligomers, the agglutination could be rather effective [77]. It is also possible that, Hsp70 occur in micelles and/or in smaller homo/heterotypic complexes also known to enhance agglutination in saliva [99]. However it is not yet known whether bacterial binding can occur under such conditions or not. The other hypothesized mechanism is based on the finding that, fungicidal activity of salivary Histatin-5 (Hst5) is initiated by binding to surface Hsp70 (Ssa1p, Ssa2p) of *Candida albicans*, followed by internalization and later to cell death [59,60]. This finding indicates that, Histatin-5 may bind salivary Hsp70 too. Although there is no evidence that, such expected salivary Hsp70/Histatin-5 complex would be able to enter and destroy *C. albicans*, but this possibility should rather be considered than simply excluded.

SALIVARY HSP70 AND ORAL DISORDERS

Oral Ulcerations

The role of Hsp70 in the formation and healing of oral ulcers was not yet studied in detail, but possible defense function of Hsp70 against oral ulceration may be expected based on the data related to gastric ulcers. In case of gastric ulcers Hsp70 is markedly overexpressed in cells located at the ulcer base [105], the level decreases with ulcer healing [105], and the extent of Hsp70 induction in mucosal cells is inversely correlates with the severity of newly induced ulcers [95,98]. These data indicate important defense role of intracellular Hsp70 against gastric ulcers. Similar intracellular response against ulcer formation may also be expected in the case of oral mucosa, but in the absence of evidence it remains a hypothesis only.

Beside intracellular defense, a coupled extracellular defense is also likely in the case of gastric ulcers, as indicated by two important findings. In one respect the regulation of gastric intracellular Hsp70 level is mediated by alpha 1A-adrenoceptors [95] the same receptor type which seems to be responsible for extracellular release (increased blood level) of Hsp70 [50]. On the other hand forced intracellular expression of Hsp70 seems to increase the secretion of Hsp70 into the extracellular space [111]. It is not yet known whether such coupling of intra and extracellular surface defense would exist in case of oral ulcers or not. But the possibility of extracellular defense in case of oral ulcers is at least given by the fact that, salivary Hsp70 is present on the ulcer surfaces covered by saliva.

There are three facets of the expected functions of salivary Hsp70 related to oral ulcers [29]. One of them can be recognized as the regular cytoprotective effect of extracellular Hsp70 as detailed above. The second one is based on the protection and repair (chaperoning) [23,24] of other salivary defense proteins (i.e. sIgA, lysozyme, cystatins, histatins, amylase, etc. [28]). Such repair activity could be especially important in the case of sIgA during

induction of antibody catalyzed ozone formation on the surface of neutrophil PMN cells [76,114]. The third facet is based on the known immunological functions of extracellular Hsp70 (see detailed above). These immune functions could be rather effective in this case, because inflammatory serum exudate containing a high amount of complement system, immunoglobulin, immune/inflammatory mediators, PMN leukocytes, and monocytes/macrophages are usually also present on the surface of oral ulcers [29].

Oral Wound Healing

Although oral tissues were not particularly investigated, based on skin experiments the effect of Hsp70 seems to be rather similar to that of ulcers also in the case of wounds. The level of intracellular Hsp70 positively correlates with the efficiency of healing [67,82], and the level of Hsp70 decreases with the progress of the healing process [67] similarly to ulcers. A coupling of intra- and extracellular Hsp70 response is also likely in this case, because of similar considerations explained in connection with ulcers (see above). The coupling of intra- and extracellular response in case of wound healing is also established by the findings that, extracellular Hsp70 is released by white blood cells in wound fluid [7]; and *in vivo* delivery of Hsp70 increased the efficiency of wound healing by the stimulation of macrophage-mediated phagocytosis of wound debris [54].

Gingival Inflammation

Gingivitis is an inflammation indicative of poor oral hygiene, without destruction of the periodontal tissues [84]. Data in the literature indicate that, the strength of the local immune reaction against microorganisms is crucial to prevent the passing of this condition to periodontitis which is a more severe inflammation with irreversible destruction of the periodontal tissues [62,84]. In case of gingivitis an inflammatory exudate appears in the periodontal sulcus containing molecular constituents such as complement system, immunoglobulin, immune and inflammatory mediators, and immune active cells like PMN leukocytes and monocytes/macrophages [29]. Since there is no pocket formation, salivary molecules are blended with the majority of the inflammatory exudate (gingival crevicular fluid) at the marginal region of the gingival sulcus. Consequently, expected immunological functions of salivary Hsp70 (see above) come into full display because of direct contact with the immune system [29].

Although the role of salivary Hsp70 is likely protective against gingivitis in majority, there are two expected mechanisms by which salivary Hsp70 may induce gingivitis under certain conditions. One of these mechanisms will be discussed later in connection with allergic and autoimmune disorders. The other one is based on the fact that, Hsp70 is secreted mainly in a lipid bound form (i.e. exosomes, lipid rafts). Protein bound lipid molecules are known to form nucleus of calcium-phosphate deposition leading to calculus formation especially in higher pH [28,48]. Since salivary Hsp70 takes part in the acquired pellicle

formation on tooth surfaces (see above), it may play a role in calculus formation leading to gingival inflammation under certain conditions.

Irritation Caused by Dentures

The most frequent reason of denture induced irritation is given by appearance of small mucosal lesions, induced by wearing removable dentures. The larger lesions are usually easily recognized, and will be healed following the correction of the denture. However the small ones (the "micro lesions") are not easily detected. Such lesion may lead to irritative-allergic reactions (see below); because of the appearance of several immune cells and immune molecules on the lesion's surface [29]. Similarly to ulcer- and wound healing (see above) the cytoprotective and immunological effects of salivary Hsp70 may be resulted in advantageously shorter healing time of such lesions. Another effect may be resulted from the fact that, salivary Hsp70 binds acrylic resins (authors unpublished data), which are basic components of removable dentures' base plates. Consequently salivary Hsp70 is likely present on the surface of dentures as constitute of acquired pellicle of dentures. Acquired pellicle of denture surfaces is a major determinant of interaction between denture and fluid-film of saliva between dentures and mucosa. This interaction is rather important in maintaining stability of removable dentures (i.e. "gluing effect" of saliva) especially in full denture cases. Because of the known protein-binding property of Hsp70, it is likely that, salivary Hsp70 plays a role in decreasing mechanical surface irritation by improving the stability of removable dentures.

Allergic and Autoimmune Reactions

There is increasing evidence that, antibodies against human Hsp of the 60 kDa family may serve as autoantigens and could initiate an autoimmune response that contributes to the initiation of gingivitis [96]. Since antibodies against salivary Hsp70 [86] likely also present, it may not be excluded that, salivary Hsp70 also induces such autoimmune mechanisms (although it should be noted that, antibodies against different heat shock proteins are differing from each other in their antigen specificity and complement activating ability [89]). Some data in the literature also suggest that, overexpression of Hsp70 may play a role in the appearance of atopic-type (IgE mediated) allergic reactions [40,61], in autoimmune disorders [91,115], and in haptenation of peptides inducing T cell immunity and sensitization [64]. However it is not yet clear to what extent extracellular Hsp70s are involved in such processes.

Evaluating the role of salivary Hsp70 in allergic and autoimmune reactions it should also be considered that, Hsp70 exerts immune regulatory and anti-inflammatory functions too [85,104,108]. Administration of recombinant Hsp70 may also resulted in attenuation of experimental autoimmune diseases [53,113]. It was also shown that, low-affinity T cells are reactive against autologous heat shock proteins [74], which might lead to generation of Th2 (IL-4- and IL-10-producing), Th3 (transforming growth factor- β -producing), or Tr1 (IL-10-

producing) regulatory T cell responses and consequent release of premised regulatory cytokines [85,104,108].

Taking together these data it is likely that, Hsp70 may take part in both the release and the control of the above allergic and autoimmune reactions under certain conditions. The basis of such "Janus-faced" behavior of Hsp70s may be at least partially rooted in that, these molecules possess both stimulating and inhibitory epitopes toward cytokine production and maturation of dendritic cells [112]. Since the oral mucosa is extensively populated by dendritic cells [25] such epitopes of salivary Hsp70 may play key role in both initiation and regulation of oral mucosal immune reactions including allergic and autoimmune pathologies.

Tumor Formation

As mentioned above a major facet of immune activation caused by Hsp70 involves the appearance of Hsp70 on the surface of certain tumor cells leading to lysis of them by natural killer (NK) cells [11,72,73]. Since in this case the tumor cell expresses its own Hsp70 on the surface, this facet is not expected for salivary Hsp70 in general. However, new findings also indicate that, surface expression of Hsp70 on tumor cells is accompanied by Hsp70 release (via lipid rafts and/or exosomes) leading to activation of NK cells also in the environment [4,6,38]. Similarly, Langerhans and dendritic cells may also be targets of such signaling [18,112], leading to increased uptake and cross presentation of tumor derived Hsp70-peptide complexes for T cell recognition [18,81].

Since there is a high number of Langerhans and dendritic cells present in superficial layer of oral mucosa [25], salivary Hsp70 may act as a confluent stimulator of the immune surveillance of these cells [29], strengthening the effectiveness (increase the perceptibility) of the premised danger signaling of early stage in situ mucosal carcinomas. Since the expression of Hsp70 seems to be positively correlated with the malignancy of oral squamous cell carcinomas [52], this effect may increase with the malignancy of the tumor.

REGULATION OF SALIVARY HSP70 LEVEL

Alteration under Psychological Stress Condition

Increase of extracellular Hsp70 level in blood [35] and human saliva [31] in response to psychological stress were demonstrated in previous studies [31,35]. In relation to *blood* a hepatosplanchnic source [32] and also brain [57] is known to be able to release high amount of Hsp70, but it is not yet known if these sources are involved in the elevation of blood level in this case. The increase of Hsp70 in the blood seems to be regulated through α 1-adrenergic receptors of not yet known target cells [35,50]. An activation of the sympathetic nervous system and consequent release of norepinephrine is likely one of the major activation pathway [35,50]. Sympathetic activation seems to be rather important also in the regulation of *salivary* Hsp70 level under stress conditions. Since α 1-adrenergic receptors seem to be widespread in both acinar and ductal cells of human submandibular glands [46], it is possible

that, α 1-adrenerg stimulation induces secretion of salivary Hsp70 in salivary glands. Further, combined α - plus β -adrenergic stimulation increases permeability of salivary glands to blood molecules in animal model [37]. Such mechanism could also be a reason of increased salivary Hsp70 level under stress conditions accompanied by increased blood Hsp70 level. Finally, elevation of salivary Hsp70 concentration and output values were found to be accompanied by elevation of salivary amylase *concentration* [31]. This finding also indicate strong connections between psychological stress, sympathetic activation and regulation of salivary Hsp70 level, because increased amylase *concentration* is a known indicator of both psychological stress [75] and stimulation of salivary glands via sympathetic nerve [88] and/or circulating catecholamine [1].

Stimulation with Chewing and Taste

Secretion of saliva is dependent upon stimuli from autonomic nerves that are the most important effector arms of reflexes activated by taste and chewing stimuli [87]. Release of proteins can be initiated by both sympathetic and parasympathetic stimuli [1,88]. Animal studies indicated that, fluid secretion is predominantly evoked by parasympathetic nerve-mediated stimuli connected with moderate protein release, whereas sympathetic nerve impulses evoke little fluid secretion with a very high amount of released secretory proteins (i.e. amylase) [1,88]. Using both chewing and taste stimuli together (i.e. mint taste chewing gum) a roughly eight fold increase of secretory rate with nearly constant protein concentration occurred indicating a predominantly parasympathetic character of the stimuli [30]. Under this condition the total Hsp70 output into the saliva increased roughly three times indicating that release of salivary Hsp70 can be stimulated also in this way [30].

Stimulation with Physiotherapy (Massage, Local Heat)

There is high number of stimulations (including mechanical stress and heat) known to induce increased expression of Hsp70 at cellular level. It is also likely that, forced intracellular expression of Hsp70 may increase the secretion of Hsp70 into the extracellular space [111]. Consequently, it is likely that, physiotherapeutic procedures like massage (mechanical stress) and local heat stimulation of major salivary glands may induce Hsp70 release into the saliva. This hypothesis was tested in a pilot study using massage and local heat in the parotid- and submandibular regions [33]. Results indicated that, total output and concentration values of salivary Hsp70 increase in both cases. Increase occurred immediately in both case, however highest values occurred initially in case of massage, whereas with a delay in case of heat stimulation [33]. Above data indicate that, salivary Hsp70 is inducible via such stimulations, however it is not jet clear if local vasodilatation, cellular damage or nervous regulatory pathways of the glands are responsible for the effect [33].

CONCLUSION

Salivary Hsp70 is likely an important factor of the oral defense mechanisms in both healthy and pathological conditions. However our knowledge related to salivary Hsp70 is still rather limited. Consequently the expectations about several functions detailed above are based on data related to extracellular but not particularly salivary Hsp70. The possible effects of special oral environment (including other salivary molecules and certain specificities of the oral mucosa) on the extracellular function of Hsp70 are not yet known. Besides summarizing the possible role of salivary Hsp70 in the oral cavity (still remaining roughly a hypothesis) another goal of the present review was to stimulate research activity leading to clear evidence related to functions of salivary Hsp70.

REFERENCES

- [1] Anderson, LC; Garrett, JR; Johnson, DA; Kauffman, DL; Keller, PJ; Thulin, A. Influence of circulating catecholamine on protein secretion into rat parotid saliva during parasympathetic stimulation. *J Physiol*, 1984 352, 163-171.
- [2] Asea, A; Kraeft, SK; Kurt-Jones, EA; Stevenson, MA; Chen, LB; Finberg, RW; Koo GC; Calderwood, SK. Hsp70 stimulates cytokine production through a CD14-dependent pathway, demonstrating its dual role as a chaperone and cytokine. *Nat Med*, 2000 6, 435-442.
- [3] Asea, A; Rehli, M; Kabingu, E; Boch, JA; Bare, O; Auron, PE; Stevenson MA; Calderwood, SK. Novel signal transduction pathway utilized by extracellular HSP70: role of toll like receptor (TLR) 2 and TLR4. *J Biol Chem*, 2002 277, 15028-15034.
- [4] Barreto, A; Gonzalez, J; Kabingu, E; Asea, A; Fiorentino, S. Stress-induced release of HSC70 from human tumors. *Cell Immunol*, 2003 222, 97-104.
- [5] Basu, S; Binder, RJ; Ramalingam, T; Srivastava, PK. CD91 is a common receptor for heat shock proteins gp96, hsp90, hsp70, and calreticulin. *Immunity*, 2001 14, 303-313.
- [6] Bausero, MA; Gastpar, R; Multhoff, G; Asea, A. Alternative mechanism by which IFN-gamma enhances tumor recognition: active release of heat shock protein 72. *J Immunol*, 2005 175, 2900-2912.
- [7] Becker, ST; Wendel, HP; Vonthein, R; Rennekampff, HO; Aebert, H. Inducible Hsp70 levels in thoracic wound fluid indicate myocardial damage after open heart surgery. *J Cardiovasc Surg (Torino)*, 2007 48, 233-237.
- [8] Becker, T; Hartl, FU; Wieland, F. CD40, an extracellular receptor for binding and uptake of Hsp70-peptide complexes. *J Cell Biol*, 2002 158, 1277-1285.
- [9] Berkovitz, BKB; Holland, GR; Moxham, BJ. *A colour atlas & textbook of oral anatomy*. (3rd impression). London: Wolfe Medical Publications Ltd.; 1984. 124-136.
- [10] Blachere, NE; Li, Z; Chandawarkar, RY; Suto, R; Jaikaria, NS; Basu, S; Udono, H; Srivastava, PK. Heat shock protein-peptide complexes, reconstituted *in vitro*, elicit peptide specific cytotoxic T-lymphocyte response and tumor immunity. *J Exp Med*, 1997 186, 1315-1322.

- [11] Botzler, C; Kolb, HJ; Issels, RD; Multhoff G. Noncytotoxic alkyl-lysophospholipid treatment increases sensitivity of leukemic K562 cells to lysis by natural killer (NK) cells. *Int J Cancer*, 1996 65, 633-638.
- [12] Boulanger, J; Faulds, D; Eddy, EM; Lingwood, CA. Members of the 70 kDa heat shock protein family specifically recognize sulfoglycolipids: role in gameterecognition and mycoplasma-related infertility. *J Cell Physiol*, 1995 165; 7-17.
- [13] Breloer, M; Fleshner, B; Bonin, AV. *In vivo* and *in vitro* activation of T cells after administration of Ag-negative heat shock proteins. *J Immunol*, 1999 162; 3141-3147.
- [14] Broquet, AH; Thomas, G; Masliah, J; Trugnan, G; Bachelet M. Expression of the molecular chaperone Hsp70 in detergent-resistant microdomains correlates with its membrane delivery and release. *J Biol Chem*, 2003 278, 21601-21606.
- [15] Buddecke, E. *Biochemische Grundlagen der Zahnmedizin*. Berlin, New York: deGruyter; 1981; 86-104.
- [16] Calderwood, SK. Evolving connections between molecular chaperones and neuronal function. *Int J Hypertherm*, 2005 21, 375-378.
- [17] Calderwood, SK; Mambula, SS; Gray jr., PJ; Theriault, JR. Extracellular heat shock proteins in cell signaling. (Minireview). *FEBS Letters*, 2007 581, 3689-3694.
- [18] Calderwood, SK; Theriault, JR; Gong, J. Message in a bottle: role of the 70-kDa heat shock protein family in anti-tumor immunity. *Eur J Immunol*, 2005 35, 2518-2527.
- [19] Campisi, J; Leem, TH; Fleshner, M. Stress-induced extracellular Hsp72 is a functionally significant danger signal to the immune system. *Cell Stress Chaperones*, 2003 8, 272-286.
- [20] Chen, S; Bawa, D; Besshoh, S; Gurd, JW; Brown IR. Association of heat shock proteins and neuronal membrane components with lipid rafts from the rat brain. *J Neurosci Res*, 2005 81, 522-529.
- [21] Chen, S; Brown, IR. Translocation of constitutively expressed heat shock protein Hsc70 to synapse-enriched areas of the cerebral cortex after hyperthermic stress. *J Neurosci Res*, 2007 85, 402-409.
- [22] Clayton, A; Turkes, A; Navabi, H; Mason, MD; Tabi, Z. Induction of heat shock proteins in B-cell exosomes. *J Cell Sci*, 2005 118, 3631-3638.
- [23] Csermely, P. Proteins, RNA-s and chaperones in enzyme evolution: a folding perspective. *Trends Biochem Sci*, 1997 22, 147-149.
- [24] Csermely, P. Chaperone-percolator model: a possible molecular mechanism of Anfinsen-cage-type chaperones. *BioEssay*, 1999 21, 959-965.
- [25] Cutler, CW; Jotwani R. Dendritic cells at the oral mucosal interface. *J Dent Res*, 2006 85, 678-689.
- [26] Daugaard, M; Rohde, M; Jäättelä M. The heat shock protein family: Highly homologous proteins with overlapping and distinct functions. (Minireview). *FEBS Letters*, 2007 581, 3702-3710.
- [27] Deocaris, CC; Kaul, SC; Wadhwa, R. On the brotherhood of the mitochondrial chaperones mortalin and heat shock protein 60. *Cell Stress Chaperon*, 2006 11, 116-128.

- [28] Fábián, TK; Fejérdy, P; Csermely, P. Chemical biology of saliva in health and disease. In: Begley T, editor. *Wiley Encyclopedia of Chemical Biology*. Hoboken: John Wiley & Sons; in press.
- [29] Fábián, TK; Fejérdy, P; Nguyen, MT; Söti, Cs; Csermely, P. Potential immunological functions of the salivary chaperone, Hsp70 in the mucosal and periodontal defense mechanisms (Review). *Arch Immunol Ther Exp*, 2007 55, 91-98.
- [30] Fábián, TK; Gáspár, J; Fejérdy, L; Kaán, B; Bálint, M; Csermely, P; Fejérdy, P. Hsp-70 is present in human saliva. *Med Sci Monit*, 2003 9, BR62-65.
- [31] Fábián, TK; Tóth, Zs; Fejérdy, L; Kaán, B; Csermely, P; Fejérdy, P. Photo-acoustic stimulation increases the amount of 70 kDa heat shock protein (Hsp70) in human whole saliva. A pilot study. *Int J Psychophysiol*, 2004 52, 211-216.
- [32] Febbraio, MA; Ott, P; Nielsen, HB; Steensberg, A; Keller, C; Krstrup, P; Secher, NH; Pedersen, BK. Exercise induces hepatosplanchnic release of heat shock protein 72 in humans. *J Physiol*, 2002 544, 957-962.
- [33] Fejérdy, L; Tóth, Zs; Kaán, B; Fábián, TK; Csermely, P; Fejérdy, P. Lokális hő- és mechanikai stressz (masszázs) stimuláció hatása az emberi kevert nyál molekuláris chaperon (Hsp70) koncentrációjára. Előzetes Vizsgálatok. *Fogorvosi Szemle*, 2004 97, 204-210.
- [34] Fink, A. Chaperone mediated protein folding. *Physiol Rev*, 1999 79, 425-449.
- [35] Fleshner, M; Campisi, J; Amiri, L; Diamond, DM. Cat exposure induces both intra- and extracellular Hsp72: the role of adrenal hormones. *Psychoneuroendocrinology*, 2004 29, 1142-1152.
- [36] Floto, RA; MacAry, PA; Boname, JM; Mien, TS; Kampmann, B; Hair, JR; Huey, OS; Houben, ENG; Pieters, J; Day, C; Oehlmann, W; Singh, M; Smith, KGC; Lehner, PJ. Dendritic cell stimulation by mycobacterial Hsp70 is mediated through CCR5. *Science*, 2006 314, 454-458.
- [37] Garrett, JR. Permeability of salivary glands to macromolecules. In: Zelles T, editor. *Advances in Physiological Sciences Vol.28.: Saliva and salivation*. Budapest: Pergamon Press - Akadémiai Kiadó; 1981; 109-118.
- [38] Gastpar, R; Gehrmann, M; Bausero, MA; Asea, A; Gross, C; Schroeder, JA; Multhoff, G. Heat shock protein 70 surface-positive tumor exosomes stimulate migratory and cytolytic activity of natural killer cells. *Cancer Res*, 2005 65, 5238-5247.
- [39] Gething, MJ. Role and regulation of the ER chaperone BiP. *Semin Cell Dev Biol*, 1999 10, 465-472.
- [40] Ghoreishi, M; Yokozeki, H; Hua, WM; Nishioka, K. Expression of 27 KD, 65 KD and 72/73 KD heat shock protein in atopic dermatitis. *J Dermatol*, 2000 27, 370-379.
- [41] Gunter, E; Walter, L. Genetic aspects of the Hsp70 multigene family in vertebrates. *Experientia*, 1994 50, 987-1001.
- [42] Guzhova, IV; Arnholdt, AC; Darieva, ZA; Kinev, AV; Launskaja, EB; Nilsson, K; Bozhkov, VM; Voronin, AP; Margulis, BA. Effects of exogenous stress protein 70 on the functional properties of human promonocytes through binding to cell surface and internalization. *Cell Stress Chaperones*, 1998 3, 67-77.

- [43] Guzhova, I; Kislyakova, K; Moskaliova, O; Fridlanskaya, I; Tytell, M; Cheetham, M; Margulis B. *In vitro* studies show that, Hsp70 can be released by glia and that exogenous Hsp70 can enhance neuronal stress tolerance. *Brain Res*, 2001 914, 66-73.
- [44] Gyrd-Hansen, M; Nylandsted, J; Jäättelä M. Heat shock protein 70 promotes cancer cell viability by safeguarding lysosomal integrity. *Cell Cycle*, 2004 3, 1484-1485.
- [45] Hartl, U. Molecular chaperones in cellular protein folding. *Nature*, 1996 381, 571-580.
- [46] Huang, Z.; Wu, L.L.; Zhang, Y.Y.; Gao, Y.; Yu, G.Y. Functional α_1 -adrenoceptor subtypes in human submandibular glands. *J Dent Res*, 2006 85(3), 251-256.
- [47] Ito, H; Takekoshi, T; Miyauchi, M; Ogawa, I; Takata, T; Nikai, H. Takemoto, K. Three-dimensional appearance of Langerhans cells in human gingival epithelium as revealed by confocal laser scanning microscopy. *Arch Oral Biol*, 1998 43, 741-744.
- [48] Jin, Y; Yip, HK. Supragingival calculus: formation and control. *Crit Rev Oral Biol Med*, 2002 13, 426-441.
- [49] Johnson, AD; Tytell, M. Exogenous Hsp70 becomes cell associated, but not internalized by stressed arterial smooth muscle cells. *In Vitro Cell Dev Biol Anim*, 1993 29, 807-812.
- [50] Johnson, JD; Fleshner, M. Releasing signals, secretory pathways, and immune function of endogenous extracellular heat shock protein 72. *J Leukoc Biol*, 2006 79, 425-434.
- [51] Kaufman, E; Lamster, IB. The diagnostic application of saliva - a review. *Crit Rev Oral Biol Med*, 2002 13, 197-202.
- [52] Kaur, J; Srivastava, A; Ralhan, R. Expression of 70-kDa heat shock protein in oral lesions: marker of biological stress or pathogenicity. *Oral Oncology*, 1998 34, 496-501.
- [53] Kingston, AE; Hicks, CA; Colston, MJ; Billingham, ME. A 71 kDa heat shock protein (hsp) from *Mycobacterium tuberculosis* has modulatory effects on experimental rat arthritis. *Clin Exp Immunol*, 1996 103, 77-82.
- [54] Kovalchin, JT; Wang, R; Wagh, MS; Azoulay, J; Sanders, M; Chandawarkar, RY. In vivo delivery of heat shock protein 70 accelerates wound healing by up-regulating macrophage-mediated phagocytosis. *Wound Repair Regen*, 2006 14, 129-137.
- [55] Lancaster, GI; Febbraio, MA. Exosome-dependent trafficking of HSP70. *J Biol Chem*, 2005 280, 23349-23355.
- [56] Lancaster, GI; Febbraio, MA. Mechanisms of stress-induced cellular HSP72 release: implications for exercise-induced increases in extracellular HSP72. *Exerc Immunol Rev*, 2005 11, 46-52.
- [57] Lancaster, GI; Moller, K; Nielsen, B; Secher, NH; Febbraio, MA; Nybo, L. Exercise induces the release of heat shock protein 72 from the human brain *in vivo*. *Cell Stress Chaperones*, 2004 9, 276-280.
- [58] Law, JS; Nelson, N; Watanabe, K; Henkin, RI. Human salivary gustin is a potent activator of calmodulin-dependent brain phosphodiesterase. *Proc Natl Acad Sci*, 1987 84, 1674-1678.
- [59] Li, XS; Sun, JN; Okamoto-Shibayama, K; Edgerton, M. *Candida albicans* cell wall Ssa proteins bind and facilitate import of salivary Histatin 5 required for toxicity. *J Biol Chem*, 2006 281, 22453-22463.

- [60] Li, XS; Reddy, MS; Baev, D; Edgerton, M. *Candida albicans* Ssa1/2p is the cell envelope binding protein for human salivary histatin 5. *J Biol Chem*, 2003 278, 28553-28561.
- [61] Liu, L; Xiao, C; Zhang, M; Cheng, L; Wang, E; Wu, T. Expression of HSP70 in peripheral lymphocytes of the patients with allergic rhinitis. *J Huazhong Univ Sci Technology Med Sci*, 2003 23, 310-312.
- [62] Lopatin, DE; Shelburne CE; Van Poperin, N; Kowalski, CJ; Bagramian, RA. Humoral immunity to stress proteins and periodontal disease. *J Periodontol*, 1999 70, 1185-1193.
- [63] Mambula, SS; Calderwood, SK. Heat shock protein 70 is secreted from tumor cells by a nonclassical pathway involving lysosomal endosomes. *J Immunol*, 2006 177, 7849-7857.
- [64] Martin, AM; Nolan, D; Gaudieri, S; Almelda, CA; Nolan, R; James, I; Carvahlo, F; Philips, E; Christiansen, FT; Purcell, AW; McCluskey, J; Mallal, S. Predisposition to abacavir hypersensitivity conferred by HLA-B*5701 and a haplotypic Hsp70-hom variant. *Proc Natl Acad Sci USA*, 2004 101, 4180-4185.
- [65] Matzinger, P. Tolerance, danger, and the extended family. *Annu Rev Immunol*, 1994 12, 991-1045.
- [66] Matzinger, P. The danger model: a renewed sense of self. *Science*, 296, 301-305.
- [67] McMurtry, AL; Cho, K; Young, LJT; Nelson, CF; Greenhalg, DG. Expression of Hsp70 in healing wounds of diabetic and nondiabetic mice. *J Surgical Research*, 1999 86, 36-41.
- [68] Milani, V; Noesser, E; Ghose, S; Kuppner, M; Ahrens, B; Scharner, A; Gastpar, R; Issels, RD. Heat shock protein 70: role in antigen presentation and immune stimulation. *Int J Hyperthermia*, 2002 18, 563-575.
- [69] Moseley P. Stress proteins and the immune response. *Immunopharmacology*, 2000 48, 299-302.
- [70] Mosser, DD; Caron, AW; Bourget, L; Meriin, AB; Sherman, MY; Morimoto, RI; Massie, B. The chaperone function of Hsp70 is required for protection against stress-induced apoptosis. *Mol Cell Biol*, 2000 20, 7146-7159.
- [71] Multhoff, G. Activation of natural killer cells by heat shock protein 70. *Int J Hyperthermia*, 2002 18, 576-585.
- [72] Multhoff, G; Botzler, C; Jennen, L; Schmidt, J; Ellwart, J; Issels, R. Heat shock protein 72 on tumor cells: a recognition structure for natural killer cells. *J Immunol*, 1997 158, 4341-4350.
- [73] Multhoff, G; Botzler, C; Wiesnet, M; Eissner, G; Issels, R. CD3⁺ large granular lymphocytes recognize a heat-inducible immunogenic determinant associated with the 72 kD heat shock protein on human sarcoma cells. *Blood*, 1995 86, 1374-1382.
- [74] Munk, ME; Schoel, B; Modrow, S; Karr, RW; Young, RA; Kaufmann, SH. T lymphocytes from healthy individuals with specificity to self-epitopes shared by the mycobacterial and human 65-kilodalton heat shock protein. *J Immunol*, 1989 143, 2844-2849.

- [75] Nater, UM; Rohleder, N; Gaab, J; Simona, B; Andreas, J; Kirschbaum, C; Ehlert, U. Human salivary alpha-amylase reactivity in a psychosocial stress paradigm. *Int J Psychophysiol*, 2005 55, 333-342.
- [76] Nathan, C. Catalytic antibody bridges innate and adaptive immunity. *Science*, 2002 298, 2143-2144.
- [77] Nemoto, TK; Fukuma, Y; Itoh, H; Takagi, T; Ono, T. A disulfide bridge mediated by cysteine 574 is formed in the dimer of the 70-kDa heat shock protein. *J Biochem (Tokyo)*, 2006 139, 677-687.
- [78] Nguyen, MT; Fábán, TK; Singh, M; Csermely, P; Söti Cs. Bacterial binding and opsonizing effect of extracellular Hsp70. Budapest Meeting Abstracts 2007. *Cell Stress & Chaperones online* 12: 1D-06-P. (www.bioone.org).
- [79] Ni, M; Lee, AS. ER chaperones in mammalian development and human diseases. (Minireview). *FEBS Letters*, 2007 581, 3641-3651.
- [80] Nollen, E; Morimoto RI. Chaperoning signaling pathways: molecular chaperones as stress-sensing "heat shock" proteins. *J Cell Sci*, 2002 115, 2809-2816.
- [81] Noessner, E; Gastpar, R; Milani, V; Brandl, A; Hutzler, PJS; Kuppner, MC; Roos, M; Kremmer, E; Asea, A. Tumor-derived heat shock protein 70 peptide complexes are cross-presented by human dendritic cells. *J Immunol*, 2002 169, 5424-5432.
- [82] Oberringer, M; Baum, HP; Jung, V; Welter, C; Frank, J; Kuhlmann, M; Mutschler, W; Hanselmann, RG. Differential expression of heat shock protein 70 in well healing and chronic human wound tissue. *Biochem Biophys Res Commun*, 1995 214, 1009-1014.
- [83] Panjwani, NN; Popova, L; Srivastava, PK. Heat shock proteins gp96 and Hsp70 activate the release of nitric oxide by APCs. *J Immunol*, 2002 168, 2997-3003.
- [84] Petit, MDA; Wassenaar, A; van der Velden, U; van Eden, W; Loos, BG. Depressed responsiveness of peripheral blood mononuclear cells to heat-shock proteins in periodontitis patients. *J Dent Res*, 1999 78, 1393-1400.
- [85] Pockley, AG. Heat shock proteins as regulators of the immune response. *Lancet*, 2003 362, 469-476.
- [86] Pockley, AG; Shepherd, J; Corton, JM. Detection of heat shock protein 70 (Hsp70) and anti-Hsp70 antibodies in the serum of normal individuals. *Immunol Investig*, 1998 27, 367-377.
- [87] Proctor, GB; Carpenter, GH. Chewing stimulates secretion of human salivary secretory immunoglobulin A. *J Dent Res*, 2001 80, 909-913.
- [88] Proctor, GB; Carpenter, GH; Anderson, LC; Garrett, JR. Nerve-evoked secretion of immunoglobulin A in relation to other proteins by parotid glands in anaesthetized rat. *Experimental Physiology*, 2000 85, 511-518.
- [89] Prohászka, Z; Duba, J; Lakos, G; Kiss, E; Varga, L; Jánoskúti, L; Császár, A; Karádi, I; Nagy, K; Singh, M; Romics, L; Füst, G. Antibodies against human heat-shock protein (hsp) 60 and micobacterial hsp65 differ in their antigen specificity and complement-activating ability. *Int Immunol*, 1999 11, 1363-1370.
- [90] Prohászka, Z; Sing, M; Nagy, K; Kiss, E; Lakos, G; Duba, J; Füst, G. Heat shock protein 70 is a potent activator of the human complement system. *Cell Stress Chaperones*, 2002 7, 17-22.

- [91] Purcell, AW; Todd, A; Kinoshita, G; Lynch, TA; Keech, CL; Gething, MJ; Gordon, TP. Association of stress proteins with autoantigens: a possible mechanism for triggering autoimmunity? *Clin Exp Immunol*, 2003 132, 193-200.
- [92] Raulston, JE; Davis, CH; Schmiel, DH; Morgan, MW; Wyrick PB. Molecular characterization and outer membrane association of a *Chlamidia trachomatis* protein related to the hsp70 family of proteins. *J Biol Chem*, 1993 268, 23139-23147.
- [93] Rea, IM; McNerlan, S; Pockley, AG. Serum heat shock protein and anti-heat shock protein antibody levels in aging. *Exp Gerontol*, 2001 36, 341-352.
- [94] Rohde, M; Dugaard, M; Jensen, MH; Helin, K; Nylanstedt, J; Jäättelä M. Members of heat-shock protein 70 family promote cancer cell growth by distinct mechanisms. *Genes Dev*, 2005 19, 570-582.
- [95] Rokutan, K. Role of heat shock proteins in gastric mucosal protection. *J Gastroenterol Hepathol*, 2000 15, Suppl:D12-19.
- [96] Schett, G; Metzler, B; Kleindienst, R; Moschén, I; Hattmannsdorfer, R; Wolf, H; Ottenhoff, T; Xu, Q; Wick, G. Salivary anti-hsp65 antibodies as a diagnostic marker for gingivitis and a possible link to atherosclerosis. *Int Arch Allergy Immunol*, 1997 114, 246-250.
- [97] Sherman, M; Gabai, V; O'Callaghan, C; Yaglom J. Molecular chaperones regulate p53 and suppress senescence programs. (Minireview). *FEBS Letters*, 2007 581, 3711-3715.
- [98] Shichijo, K; Ihara, M; Matsuu, M; Ito, M; Okumura, Y; Sekine, I. Overexpression of heat shock protein 70 in stomach of stress induced gastric ulcer-resistant rats. *Dig Dis Sci*, 2003 48, 340-348.
- [99] Soares, RV; Lin, T; Siqueria, CC; Bruno, LS; Li, X; Oppenheim, FG; Offner, G; Troxler, RF. Salivary micelles: identification of complexes containing MG2, sIgA, lactoferrin, amylase, glycosylated proline-rich protein and lysozyme. *Arch Oral Biol*, 2004 49, 337-343.
- [100] Srivastava, PK. Heat shock proteins in innate and adaptive immunity. *Nat Rev Immunol*, 2002 2, 185-194.
- [101] Suto, R; Srivastava, PK. A mechanism for the specific immunogenicity of heat shock protein-chaperoned peptides. *Science*, 1995 269, 1585-1588.
- [102] Thériault, JR; Adachi, H; Calderwood, SK. Role of scavenger receptors in the binding and internalization of heat shock protein 70. *J Immunol*, 2006 177, 8604-8611.
- [103] Thériault, JR; Mambula, SS; Sawamura, T; Stevenson, MA; Calderwood, SK. Extracellular HSP70 binding to surface receptors present on antigen presenting cells and endothelial cells. *FEBS Letters*, 2005 579, 1951-1960.
- [104] Todryk, SM; Gough, MJ; Pockley, G. Facet of heat shock protein 70 show immunotherapeutic potential. *Immunology*, 2003 110, 1-9.
- [105] Tsukimi, Y; Nakai, H; Itoh, S; Amagase, K; Okabe, S. Involvement of heat shock proteins in the healing of acetic acid-induced gastric ulcers in rats. *J Physiol Pharmacol*, 2001 52, 391-406.
- [106] Tytell, M; Greenberg, SG; Lasek RJ. Heat shock like protein is transferred from glia to axon. *Brain Res*, 1986 363, 161-164.
- [107] Udono, H; Srivastava, PK. Heat shock protein 70-associated peptides elicit specific cancer immunity. *J Exp Med*, 1993 178, 1391-1396.

- [108] van Eden, W; van der Zee, R; Paul, AG; Prakken, BJ; Wendling, U; Anderton, SM; Wauben, MH. Do heat shock proteins control the balance of T-cell regulation in inflammatory diseases? *Immunol Today*, 1998 19, 303-307.
- [109] Vanmuylder, N; Evrard, L; Daelemans, P; Dourov, N. Chaperones in the parotid gland: localization of heat shock proteins in human adult salivary glands. *Cell Tissues Organs*, 2000 167, 199-205.
- [110] Velez-Granell, CS; Arias, AE; Torres-Ruiz, JA, Bendayan M. Molecular chaperones in pancreatic tissue: the presence of cpn10, cpn60, and Hsp70 in distinct compartment along the secretory pathway of the acinar cells. *J Cell Sci*, 1994 107, 539-549.
- [111] Wang, MH; Grossmann, ME; Young CYF. Forced expression of heat-shock protein 70 increases the secretion of Hsp70 and provides protection against tumor growth. *British J Cancer*, 2004 90, 926-931.
- [112] Wang, Y; Whittall, T; McGowan, E; Younson, J; Kelly, C; Bergmeier, LA; Singh, M; Lehner, T. Identification of stimulating and inhibitory epitopes within the heat shock protein 70 molecule that modulate cytokin production and maturation of dendritic cells. *J Immunol*, 2005 174, 3306-3316.
- [113] Wendling, U; Paul, L; van der Zee, R; Prakken, B; Singh, M; van Eden, W. A conserved mycobacterial heat shock protein (hsp) 70 sequence prevents adjuvant arthritis upon nasal administration and induces IL-10-producing T cells that cross-react with the mammalian self-hsp70 homologue. *J Immunol*, 2000 164, 2711-2717.
- [114] Wentworth, P; McDunn, JE; Wentworth, AD; Takeuchi, C; Nieva, J; Jones, T; Bautista, C; Ruedi, JM; Gutierrez, A; Janda, KD; Babior, BM; Eschenmoser, A; Lerner, RA. Evidence for antibody-catalyzed ozone formation in bacterial killing and inflammation. *Science*, 2002 298, 2195-2199.
- [115] Whittall, T; Wang, Y; Kelly, CG; Thompson, R; Sanderson, J; Lomer, M; Soon, SY; Bergmeier, LA; Singh, M; Lehner, T. Tumour necrosis factor- α production stimulated by heat shock protein 70 and its inhibition in circulating dendritic cells and cells eluted from mucosal tissues in Chron's disease. *Clin Exp Immunol*, 2006 143, 550-559.
- [116] Williams, PL., Warwick R. *Gray's Anatomy*. 36th Edition. Edinburgh, London, Melbourne, New York: Churchill Livingstone; 1980; 1272-1281.
- [117] Wright, BH; Corton JM; El-Nahas, AM; Wood, RF; Pockley AG. Elevated levels of circulating heat shock protein (Hsp70) in peripheral and renal vascular disease. *Heart Vessels*, 2000 15, 18-22.

Reviewed by Prof. Dr. Gábor Varga Ph.D., DSc., professor and chairman of the Department of Oral Biology, Semmelweis University, Budapest, Hungary

HEAT SHOCK PROTEINS: NEW RESEARCH

**EMMA MOREL AND CAMILLE VINCENT
EDITORS**

Nova Biomedical Books
New York

Copyright © 2008 by Nova Science Publishers, Inc.

All rights reserved. No part of this book may be reproduced, stored in a retrieval system or transmitted in any form or by any means: electronic, electrostatic, magnetic, tape, mechanical photocopying, recording or otherwise without the written permission of the Publisher.

For permission to use material from this book please contact us:

Telephone 631-231-7269; Fax 631-231-8175

Web Site: <http://www.novapublishers.com>

NOTICE TO THE READER

The Publisher has taken reasonable care in the preparation of this book, but makes no expressed or implied warranty of any kind and assumes no responsibility for any errors or omissions. No liability is assumed for incidental or consequential damages in connection with or arising out of information contained in this book. The Publisher shall not be liable for any special, consequential, or exemplary damages resulting, in whole or in part, from the readers' use of, or reliance upon, this material.

Independent verification should be sought for any data, advice or recommendations contained in this book. In addition, no responsibility is assumed by the publisher for any injury and/or damage to persons or property arising from any methods, products, instructions, ideas or otherwise contained in this publication.

This publication is designed to provide accurate and authoritative information with regard to the subject matter covered herein. It is sold with the clear understanding that the Publisher is not engaged in rendering legal or any other professional services. If legal or any other expert assistance is required, the services of a competent person should be sought. FROM A DECLARATION OF PARTICIPANTS JOINTLY ADOPTED BY A COMMITTEE OF THE AMERICAN BAR ASSOCIATION AND A COMMITTEE OF PUBLISHERS.

LIBRARY OF CONGRESS CATALOGING-IN-PUBLICATION DATA

Heat shock proteins: new research/Emma Morel and Camille Vincent(editors).
p.;cm.

Includes bibliographical references and index.

ISBN 978-1-60456-641-3(hardcover)

1. Heat shock proteins.I.Morel,Emma.II.Vincent,Camille,1960-
[DNLM:1.Heat-Shock Proteins.QU 55.6 H4.76 2008]

QP552.H43.H437 2008

572'6—dc22

2008013814

Published by Nova Science Publishers, Inc. + New York