

Molecular Chaperones in the Etiology and Therapy of Cancer*

Csaba SÓTI and Péter CSERMELY

Department of Medical Chemistry, Semmelweis University of Medicine, Budapest, Hungary

Molecular chaperones are ubiquitous, well-conserved proteins that account for 2–5 % of all cellular proteins in most cells. The present review summarizes our current knowledge about their involvement in the etiology and therapy of cancer with special emphasis on the expression of chaperones in malignant cells, their role in folding of (proto)onco-

gene products, cell cycle regulation, cell differentiation and apoptosis, development of metastasis, and their participation in the recognition of malignant cells. We also overview the importance of chaperones in hyperthermia, drug resistance, and recent approaches in chaperone-immunotherapy. (Pathology Oncology Research Vol 4, No 4, 316–321, 1998)

Key words: chaperone, heat shock protein, stress protein, metastasis, immunotherapy, drug resistance

Introduction

Molecular chaperones have been defined as “proteins that bind to and stabilize an otherwise unstable conformer of another protein – and, by controlled binding and release, facilitate its correct fate *in vivo*: be it folding, oligomeric assembly, transport to a particular subcellular compartment, or disposal by degradation”.³¹ Chaperones are ubiquitous, highly conserved proteins which probably played a major role in the evolution of modern enzymes.¹⁶ Chaperones are vital for our cells during their whole lifetime. However, they are needed even more after environmental stress, which induces protein damage. Stress (heat shock, major changes in the cellular environment after the activation of various pathogens or during the development of disease, etc.) induces the synthesis of many chaperones

which therefore are called heat-shock, or stress proteins. Chaperones play an essential role in the etiology of numerous diseases, with a rapidly increasing role in clinical practice.^{50,88,92} Lacking a settled view about their exact and specific cellular functions, chaperones are still best classified by their molecular weights. The major chaperone families are listed in *Table 1*.

Induction of molecular chaperones in malignant cells

Chaperones help damaged proteins to re-fold to their native conformation, therefore it is not surprising that they are expressed after the cell experiences various types of environmental stress. Cancer cells have an especially “stressful” life: lack of nutrients, oxygen, space limitations, and hostile environment are all important factors which acting by alone would induce a large number of chaperone proteins. We have summarized some examples of chaperone induction in *Table 2*. Generally, induced levels of chaperones help tumor cell survival. Overexpression of Hsp90- α is usually associated with poor prognosis in breast cancer.⁹³ However, in some cases such as after the overexpression of Hsp25, proliferation of malignant cells slows down.⁴⁷ Thus due to the pleiotropic effects of molecular chaperones. It is often difficult to make simplified statements on their association with poor or better prognosis in various types of cancer.¹⁵

Molecular chaperones are induced by almost all treatment protocols used to eliminate tumors. Hyperthermia

Received: August 24, 1998; *accepted:* Sept 20, 1998

Correspondence: Dr. Péter CSERMELY, Department of Medical Chemistry, Semmelweis University of Medicine, P. O. Box 260., H-1444 Budapest 8, Hungary; tel: +36-1-266-2755/4102; fax: +36-1-266-6550; e-mail: csermely@puskin.sote.hu

Abbreviations: Grp, glucose regulated protein; Hsp, heat shock protein.

*Research in the authors' lab was supported by grants from the Hungarian National Science Foundation (OTKA T25206), from the Hungarian Ministry of Social Welfare (ETT 493/96), from the Hungarian Ministry of Culture and Education (FKFP 761/97) and from the ICGEB. P.C. is an International Research Scholar of the Howard Hughes Medical Institute (HHMI 75195-541701).

Table 1. The major molecular chaperone families

<i>Some common names of eukaryotic chaperone family members</i>	<i>References for recent reviews</i>
Hsp25, Hsp27, crystallins, Small heat-shock proteins	10,30
Hsp60, chaperonins	11,31
Hsp70, Grp78, BiP	11,31
Hsp90, Grp94	10,17,63,64
Hsp110	68,91

Neither the co-chaperones (chaperones which help the function of other chaperones listed, such as Hsp10, dnaJ-homologues, Hip, Hop, Hup, etc.), nor the so-called folding catalysts, the peptidyl-prolyl isomerases (immunophilins) and protein disulfide isomerases were included in this table, albeit almost all of these proteins also possess a "traditional" chaperone activity in their own right.

and various forms of chemotherapy are the classical examples of treatment-induced chaperone expression, as we will discuss in later sections of this review. Chaperone induction has also been observed after radiation therapy,⁶⁷ or addition photosensitizers and phototherapy.⁵⁵

Association of chaperones with (proto)oncogene products: their role in regulation of the cell cycle, cell differentiation and apoptosis

Chaperones participate in the folding of numerous protooncogene and oncogene products. These target-proteins often form stable complexes with chaperones which keep them in an "activation-competent" state. E.g.: Hsp90 is necessary for the folding of several protooncogene/oncogene protein kinases, such as members of the src and raf families.¹⁷ Chaperones also participate in the maintenance of correct conformation of other proteins involved in signal transduction, such as receptors, G-proteins and transcription factors.⁶³ Chaperones are necessary for the folding of several proteins of oncogenic viruses, such as those of the hepatitis B virus.³⁴

The expression pattern of several chaperones is cell cycle dependent. E.g.: Hsp90- α mRNA is induced at the G₁/S transition of chicken hepatoma cells.⁴³ In eukaryotic organisms folding of *de novo* synthesized proteins is usually mediated by the ribosome-complex itself. However, folding of several key members of cell cycle regulation requires the help of specific molecular chaperones. E.g.: Hsp90 is necessary for the folding of the cyclin-dependent kinase, CDK4⁷⁹ and the cyclin dependent kinase regulator, Wee.¹ Hsp70 participates in the activation of the tumor-suppressor p53 protein.³⁷ The novel Hsp90 homologue, Hsp75/TRAP-1, as well as the constitutively expressed isoform of Hsp70 associate with the tumor suppressor retinoblastoma protein,

most probably stabilizing the conformation of its dephosphorylated, thus tumor suppressive form.^{13,38}

Changes in chaperone expression are usually accompanying the differentiation of various cell types. Differentiation of embryonal carcinoma cells leads to elevated levels of small heat shock proteins which may play an important role in various signaling events leading to the differentiated state.^{72,78} On the contrary, expression of 70 and 90 kDa chaperones is lowered when the cells leave vigorous proliferation,^{3,17,33} which may reflect a decreased need for help in conformational rearrangements.

Induction, or overexpression of various chaperones generally protects host cells from apoptosis.⁶⁵ Transgenic mice expressing the inducible form of the 70 kDa molecular chaperone, develop T cell lymphoma due to the severely impaired apoptosis in T cell selection.⁷⁰ Similarly, expression of Hsp27 or Hsp70 protects tumor cells against the apoptotic effects of tumor necrosis factor- α .^{40,90} On the contrary, overexpression of Hsp90 enhances tumor necrosis factor- α induced apoptosis of U937 cells.²⁷ This may be related to the possible involvement of an Hsp90-homologue in type-1 tumor necrosis factor receptor signaling as reported by Song et al.⁷¹ Thus the involvement of chaperones in the diversion of the normal cell cycle towards

Table 2. Induction of molecular chaperones in malignant human cells

<i>Chaperone family member</i>	<i>Malignant cell type</i>	<i>References</i>
small heat shock proteins	breast cancer	14,59
	hepatoma	18
	neuroectodermal tumor	39
Hsp60	breast cancer	7
	lymphoma	21
Hsp70	endometrial carcinoma	58
	lung carcinoma	22
	melanoma	22,60
	pancreatic carcinoma	29
	renal carcinoma	66
Grp78	breast cancer	7
Hsp90	breast cancer	22,24,41,93
	endometrial carcinoma	58
	gastrointestinal cancer	20
	hepatoma	22
	leukemia	12,94
	lung carcinoma	22
	melanoma	22,60
	microcytoma	22
	ovarian cancer	54
	pancreatic carcinoma	29
Grp94	breast cancer	7,23,24,93
	colon adenocarcinoma	53
Hsp 110 ^a		

^aInduction of Hsp110 in tumors has not been reported (yet).

apoptosis most probably depends on the type of apoptotic signal. As is obvious from the above, our present knowledge about the involvement of chaperones in the cell cycle and apoptosis is rather fragmentary. However, these areas may well provide significant major advances in the understanding of chaperone function in the near future.

Involvement of chaperones in metastasis development

Induction of various molecular chaperones, such as Hsp27,⁸⁰ Hsp70,^{45,60} Hsp90,⁴¹ and the collagen-specific chaperone, Hsp47⁵⁶ was observed in several metastasis models. The putative heparanase and protease (aminopeptidase) activities of the Hsp90-homologue, Grp94, together with its frequent expression on the surface of tumor cells,^{19,28,73,76} may enable Grp94 to act as a mediator of metastasis generation. However, the testing of the putative role of Grp94 and other chaperones in promotion of metastasis formation is a task for future research.

Surface expression of chaperones, their role in antigen-presentation, and in immunorecognition of malignant cells

In the end of the eighties Hsp70, Hsp90 and Grp94 (termed gp96) were identified as tumor-specific antigens expressed on the surface of various tumor cells.^{44,76,87} Expression of molecular chaperones on the surface of malignant cells and their secretion to the extracellular fluid has also been reported by numerous other laboratories.^{4,17,22,57} Interestingly, extracellular Hsp90- α had a stimulatory effect on the growth of some lymphoid cell lines⁴⁸ and Grp78 was identified as a potential intercellular signal-transducing protein between pancreatic cancer cells.²⁵ Presently neither the molecular details of the surface attachment of molecular chaperones, nor the exact mechanism of their secretion are known.

Though differences in protein structure of various tumor-derived, surface-expressed chaperones were minor, if anything, their immunogenicity showed major differences. This apparent discrepancy led Pramod Srivastava to suggest that the chaperone-related immunogenicity resides in a great variety of peptides, which are non-covalently associated to, and "presented" by the chaperone.^{74,75} Endogenously synthesized antigenic determinants are generally presented on major histocompatibility complex (MHC) class I molecules, whereas exogenous antigens are presented by MHC class II molecules. Heat shock and glucose-regulated proteins (Hsp70, Hsp90 and Grp94) may present their bound peptides to MHC class I molecules. Under normal (non-stressed) conditions this may be a helper mechanism for loading of the MHC class I molecules in the endoplasmic reticulum. However, stress proteins may carry their immunogen peptides to MHC class I

molecules other than those of their original cells by lysis of the original cell and subsequent phagocytosis by macrophages or by direct macrophage-engulfment of the whole original cell. Since heat shock proteins are highly conserved, transfer of their peptide-load to MHC class I molecules may also occur after the lysis or phagocytosis of foreign cells with different haplotypes. Hence foreign chaperones may "disguise" their bound foreign peptides as self. Thus insertion of the nondiscriminating stress proteins to the peptide/antigen-presenting "relay" may explain the phenomenon of cross-priming, *i.e.* that not all the processing of the antigens occurs *via* the haplotype-restricted MHC class I molecules of the immunized mouse, but at least some of peptide/antigens are salvaged by the macrophages of the immunized mouse *directly* from the chaperones of the immunizing cells (having a different haplotype). During the last years the above hypothesis of Srivastava et al.⁷⁷ has been supported by several pieces of experimental evidence.^{2,83,85,86}

The involvement of chaperones in antigen-presentation also means that, in an organism developing malignant cells, the MHC non-restricted presentation of tumor antigens becomes more dominant. This mechanism increases the efficiency of immune surveillance. Tumor-derived peptide-loaded chaperones (*via* the peptide-presenting macrophage-MHC class I molecules) may prime cytotoxic lymphocytes even after the death and lysis of the original malignant cells, which extends the cytotoxic response and also makes it more efficient.⁷⁷

Hyperthermia and cancer treatment

Hyperthermia has been utilized in clinical practice as a primary treatment or as an adjuvant to radio-/chemotherapy of cancer for a long time.⁸ Hyperthermal treatment protocols are based on proper focusing of thermal damage to cause the selective injury of tumor cells using computer-assisted 3D targeting.⁸² The current review warrants for an even sharper (theoretically: all-or-non) discrimination between target and neighboring cells since improper (sub-optimal) heating of tumor cells may induce an expression of their molecular chaperones and thus may lead to an increase in their survival and metastatic potential. Induction of Hsp70 may be efficiently used to judge the extent of hyperthermia in the malignant tissue.⁵¹ On the other hand, massive induction of Hsp70 (and other heat shock proteins, such as Hsp110) may indicate the development of tumor-thermotolerance and stress-(drug)-resistance.

Chaperones and drug resistance

Heat treatment leads to increased drug resistance in many tumor cells. In agreement with these early results, in several malignant cell types the simultaneous induction of

various heat shock proteins and multidrug resistance has been observed.^{14,36,59} Administration of chemotherapeutic agents often leads to a further increase in the expression of molecular chaperones, such as Hsp25⁶ or Hsp70.³⁵ As a further evidence suggesting the chaperone-induced protection against chemotherapy high level of Hsp60 predicts poor survival of patients treated with cisplatin-containing chemotherapy protocols.⁴⁶ However, in some cases the chaperone-induction correlates with the induction of P-glycoprotein, e.g. in case of elevated Hsp90- β levels, where the chaperone has also been shown to associate directly with the multi-drug transporter.⁵ In other cases elevated chaperone levels themselves seem to induce a "cross-resistance" against various chemotherapeutic agents irrespectively from the level of multidrug transporter present.^{14,36,59} To make the situation even more complex, overexpression of P-glycoprotein has also been observed in heat-resistant hepatoma cells, which are defective in the induction of Hsp70.^{61,62}

Molecular chaperones (the cytoplasmic "foldosome", containing Hsp70, Hsp90 and numerous co-chaperones) are actively involved in the folding and activation of steroid receptors.^{17,63,64} Thus it is not surprising that they play a pivotal role in the development of steroid resistance in various forms cancer, such as breast, endometrial and prostate cancer. Since length limitations of the present review do not allow us to review this extensive field, the interested reader is referred to other recent reviews providing an excellent summary of the subject.^{26,49,89}

Molecular chaperones in cancer immunotherapy

A large part of tumor immunogenicity resides in the great variety of chaperone-associated tumor-specific peptides. Tumor-specific, chaperone-presented peptides are taken up by macrophages and presented by the macrophage MHC class I molecules. These macrophages are able to prime cytotoxic T lymphocytes for an anti-tumor attack.^{2,77,83,86} The chaperone-mediated "escape route" of cytotoxic lymphocyte priming from the restrictive self-MHC molecules has profound consequences in cancer immunotherapy. The vaccination procedure does not necessarily have to use autologous or HLA-matched cells, but a preparation of chaperone-peptide complex from the specific tumor may be used as an effective immunogen to vaccinate the same patient. This vaccination protocol may be extended to shared tumor antigens in the future which may alleviate the need for the costly and time-consuming "personalized" vaccines.^{9,32,77,84,85} As an approach of this type, if tumor cells are transfected with the mycobacterial heat shock protein, Hsp65 (which is a major common immunogen in almost all organisms), they lose their tumorigenicity,⁵² or can be used as cancer vaccines.⁶⁹

As an alternative chaperone-based immune-related therapy, suppression of the synthesis of certain chaperones, such as the Hsp70 homologue, Grp78, eliminates the tumor resistance to cell mediated cytotoxicity.^{42,81}

Conclusions

Overexpression of chaperones in tumor cells is a rather general phenomenon caused by the increased demand of accelerated cell proliferation and the harmful environment. Chaperones protect malignant cells from many of the environmental stresses and render them more resistant against apoptosis, anticancer drugs and immune-attacks. Chaperone-induction may also lead to an increase of the metastatic potential. Tumor chaperones seem to be one of the devils of tumor therapy, thus various selective methods to impair their synthesis in tumor cells have high therapeutic potential. On the other hand, chaperone-peptide vaccination may provide a unique, a very powerful tool in cancer treatment.

References

1. *Aligue R, Akhavan-Niak H and Russell P*: A role for Hsp90 in cell cycle control: Wee1 tyrosine kinase activity requires interaction with Hsp90. *EMBO J* 13:6099-6106, 1994.
2. *Arnold D, Faath S, Rammensee H, et al*: Cross-priming of minor histocompatibility antigen-specific cytotoxic T cells upon immunization with the heat shock protein gp96. *J Exp Med* 182:885-889, 1995.
3. *Beere HM, Morimoto RI and Hickman JA*: Investigations of mechanisms of drug-induced changes in gene expression: N-methylformamide-induced changes in synthesis of the M(r) 72,000 constitutive heat shock protein during commitment of HL-60 cells to granulocyte differentiation. *Cancer Res* 53:3034-3039, 1993.
4. *Berger CL, Dong Z, Hanlon D, et al*: A lymphocyte cell surface heat shock protein homologous to the endoplasmic reticulum chaperone, immunoglobulin heavy chain binding protein BIP. *Int J Cancer* 71:1077-1085, 1997.
5. *Bertram J, Palfner K, Hiddemann W, et al*: Increase of P-glycoprotein-mediated drug resistance by hsp90- β . *Anticancer Drugs* 7:838-845, 1996.
6. *Bielka H, Hoinkis G, Oesterreich S, et al*: Induction of the small stress protein, hsp25, in Ehrlich ascites carcinoma cells by anticancer drugs. *FEBS Lett* 343:165-167, 1994.
7. *Bini L, Magi B, Marzocchi B, et al*: Protein expression profiles in human breast ductal carcinoma and histologically normal tissue. *Electrophoresis* 18:2832-2841, 1997.
8. *Bisht KS and Uma-Devi P*: Hyperthermia in cancer research: current status. *Indian J Exp Biol* 34:1183-1189, 1996.
9. *Blachere NE, Udono H, Janetzki S, et al*: Heat shock protein vaccines against cancer. *J Immunother* 14:352-356, 1993.
10. *Buchner J*: Supervising the fold: functional principles of molecular chaperones. *FASEB J* 10:10-19, 1996.
11. *Bukau B and Horwich AL*: The Hsp70 and Hsp60 chaperone machines. *Cell* 92:351-366, 1998.
12. *Chant ID, Rose PE and Morris AG*: Analysis of heat-shock protein expression in myeloid leukemia cells by flow cytometry. *Br J Haematol* 90:163-168, 1995.

13. *Chen CF, Chen Y, Dai K, et al*: A new member of the hsp90 family of molecular chaperones interacts with the retinoblastoma protein during mitosis and after heat shock. *Mol Cell Biol* 16:4691-4699, 1996.
14. *Ciocca DR, Fuqua SA, Lock-Lim S, et al*: Response to human breast cancer cells to heat shock and chemotherapeutic drugs. *Cancer Res* 52:3648-3654, 1992.
15. *Conroy SE and Latchman DS*: Do heat shock proteins have a role in breast cancer? *Br J Cancer* 74:717-721, 1996.
16. *Csermely P*: Proteins, RNA-s, chaperones and enzyme evolution: a folding perspective. *Trends Biochem Sci* 22:147-149, 1997.
17. *Csermely P, Schneider T, Sóti Cs, et al*: The 90 kDa molecular chaperone family: structure, function and clinical applications. A comprehensive review. *Pharmacol Therapeutics* 79:129-168, 1998.
18. *Delhaye M, Gulbis B, Galand P, et al*: Expression of 27-kD heat-shock protein isoforms in human neoplastic and nonneoplastic liver tissues. *Hepatology* 16:382-389, 1992.
19. *De Vouge MW, Yamazaki A, Bennett SA, et al*: Immunoselection of GRP94/endoplasmic from a KNRK cell-specific lambda gt11 library using antibodies directed against a putative heparanase amino-terminal peptide. *Int J Cancer* 56:286-294, 1994.
20. *Ehrenfried JA, Herron BE, Townsend CM, et al*: Heat shock proteins are differentially expressed in human gastrointestinal cancers. *Surg Oncol* 4:197-203, 1995.
21. *Ferm MT, Soderstrom K, Jindal S, et al*: Induction of human hsp60 expression in monocytic cell lines. *Int Immunol* 4:305-311, 1992.
22. *Ferrari M, Heltai S, Zocchi MR, et al*: Unusual expression and localization of heat-shock proteins in human tumor cells. *Int J Cancer* 51:613-619, 1992.
23. *Franzen B, Linder S, Alaiya AA, et al*: Analysis of polypeptide expression in benign and malignant human breast lesions - down regulation of cytokeratins. *Br J Canc* 74:1632-1638, 1996.
24. *Franzen B, Linder S, Alaiya AA, et al*: Analysis of polypeptide expression in benign and malignant human breast lesions. *Electrophoresis* 18:582-587, 1997.
25. *Furutani M, Arai S, Mizumoto M, et al*: Identification of a neutrophil gelatinase-associated lipocalin mRNA in human pancreatic cancers using a modified signal sequence trap method. *Cancer Lett* 122:209-214, 1998.
26. *Gadducci A and Genazzani AR*: Steroid hormones in endometrial and breast cancer. Review. *Eur J Gynaecol Oncol* 18:371-378, 1997.
27. *Galea-Lauri J, Richardson AJ, Latchman DS, et al*: Increased heat shock protein 90 (hsp90) expression leads to increased apoptosis in the monoblastoid cell line U937 following induction with TNF- α and cycloheximide: a possible role in immunopathology. *J Immunol* 157:4109-4118, 1996.
28. *Graham LD*: Tumour rejection antigens of the hsp90 family (gp96) closely resemble tumour-associated heparanase enzymes. *Biochem J* 301:917-918, 1994.
29. *Gress TM, Muller-Pillasch F, Weber C, et al*: Differential expression of heat shock proteins in pancreatic carcinoma. *Cancer Res* 54:547-551, 1994.
30. *Groenen PJ, Merck KB, de Jong WW, et al*: Structure and modifications of the junior chaperone α -crystallin. From lens transparency to molecular pathology. *Eur J Biochem* 225:1-19, 1994.
31. *Hartl F-U*: Molecular chaperones in cellular protein folding. *Nature* 381:571-580, 1996.
32. *Heike M, Noll B and Meyer zum Buschenfelde KH*: Heat shock protein-peptide complexes for use in vaccines. *J Leukoc Biol* 60:153-158, 1996.
33. *Hensold JO and Housman DE*: Decreased expression of the stress protein HSP70 is an early event in murine erythroleukemic cell differentiation. *Mol Cell Biol* 8:2219-2223, 1988.
34. *Hu J, Toft DO and Seeger C*: Hepadnavirus assembly and reverse transcription require a multi-component chaperone complex which is incorporated into nucleocapsids. *EMBO J* 16:59-68, 1997.
35. *Huber SA*: Heat-shock protein induction in adriamycin and picornavirus-infected cardiocytes. *Lab Invest* 67:218-224, 1992.
36. *Huot J, Roy G, Lambert H, et al*: Co-induction of hsp27 phosphorylation and drug resistance in chinese hamster cells. *Int J Oncology* 1:31-36, 1992.
37. *Hupp TR, Meek DW, Midgley CA, et al*: Regulation of the specific DNA binding function of p53. *Cell* 71:875-886, 1992.
38. *Inoue A, Torigoe T, Sogahata K, et al*: 70-kDa heat shock cognate protein interacts directly with the N-terminal region of the retinoblastoma gene product pRb. Identification of a novel region of pRb-mediated protein interaction. *J Biol Chem* 270:22571-22576, 1995.
39. *Iwaki T, Iwaki A, Miyazono M, et al*: Preferential expression of B-crystallin in astrocytic elements of neuroectodermal tumors. *Cancer* 63:320-322, 1991.
40. *Jaattela M, Wissing D, Bauer PA, et al*: Major heat shock protein, hsp70 protects tumor cells from tumor necrosis factor cytotoxicity. *EMBO J* 11:3507-3512, 1992.
41. *Jameel A, Skilton RA, Campbell TA, et al*: Clinical and biological significance of HSP89 α in human breast cancer. *Int J Cancer* 50:409-415, 1992.
42. *Jamora C, Dennert G and Lee AS*: Inhibition of tumor progression by suppression of stress protein GRP78/BiP induction in fibrosarcoma B/C10ME. *Proc Natl Acad Sci USA* 93:7690-7694, 1996.
43. *Jerome V, Vourch C, Baulieu EE, et al*: Cell cycle regulation of the chicken hsp90 expression. *Exp Cell Res* 205:44-51, 1993.
44. *Kanno A, Sato N, Yagihashi A, et al*: Heat-, or stress-inducible transformation-associated cell surface antigen on the activated H-ras oncogene-transfected rat fibroblast. *Cancer Res* 49:6578-6582, 1989.
45. *Kato S, Hirano A, Kato M, et al*: Stress-response (heat-shock) protein 72 expression in tumors of the central nervous system: an immunohistochemical investigation. *Acta Neuropathol* 84:261-264, 1992.
46. *Kimura E, Enns RE, Thiebaut F, et al*: Regulation of HSP60 mRNA expression in a human ovarian carcinoma cell line. *Cancer Chemother Pharmacol* 32:279-285, 1993.
47. *Knauf U, Bielka H and Gaestel M*: Over-expression of the small heat-shock protein, hsp25, inhibits growth of Ehrlich ascites tumor cells. *FEBS Lett* 309:297-302, 1992.
48. *Kuroita T, Tachibana H, Ohashi H, et al*: Growth stimulating activity of heat shock protein 90 to lymphoid cell lines in serum-free medium. *Cytotechnology* 8:109-117, 1992.
49. *el Lalani N, Laniado ME and Abel PD*: Molecular and cellular biology of prostate cancer. *Cancer Metastasis Rev* 16:29-66, 1997.
50. *Latchman DS*: Heat shock proteins and human disease. *J Roy Coll Phys London* 25:295-299, 1991.
51. *Liu FF, Miller N, Levin W, et al*: The potential role of HSP70 as an indicator of response to radiation and hyperthermia treatments for recurrent breast cancer. *Int J Hyperthermia* 12:197-208, 1996.
52. *Lukacs KV, Lowrie DB, Stokes RW, et al*: Tumor cells transfected with a bacterial heat-shock gene lose tumorigenicity and induce protection against tumors. *J Exp Med* 178:343-348, 1993.
53. *Menoret A, Meflah K and Le Pendu J*: Expression of the 100-kDa glucose-regulated protein (GRP100/endoplasmic) is associated with tumorigenicity in a model of rat colon adenocarcinoma. *Int J Cancer* 56:400-405, 1994.
54. *Mileo AM, Fanuele M, Battaglia F, et al*: Selective overexpression of mRNA coding for 90 kDa stress-protein in human ovarian cancer. *Anticancer Res* 10:903-906, 1990.

55. Morgan J, Whitaker JE and Oseroff AR: GRP78 induction by calcium ionophore potentiates photodynamic therapy using the mitochondrial targeting dye victoria blue BO. *Photochem Photobiol* 67:155-164, 1998.
56. Morino M, Tsuzuki T, Ishikawa Y, et al: Specific expression of HSP47 in human tumor cell lines in vitro. *In Vivo* 11:17-21, 1997.
57. Multhoff G and Hightower LE: Cell surface expression of heat shock proteins and the immune response. *Cell Stress Chaperones* 1:167-176, 1996.
58. Nanbu K, Konishi I, Komatsu T, et al: Expression of heat shock proteins HSP70 and HSP90 in endometrial carcinomas. Correlation with clinicopathology, sex steroid receptor status, and p53 protein expression. *Cancer* 77:330-338, 1996.
59. Oesterreich S, Weng CN, Qiu M, et al: The small heat shock protein hsp27 is correlated with growth and drug resistance in human breast cancer cell lines. *Cancer Res* 53:4443-4448, 1993.
60. Pia Protti M, Heltai S, Bellone M, et al: Constitutive expression of the heat shock protein 72 kDa in human melanoma cells. *Cancer Lett* 85:211-216, 1994.
61. Purity M, Nguyen VT, Dubois MF, et al: Decreased stress inducibility of Hsp68 protein in a rat hepatoma variant clone. *Eur J Biochem* 210:793-800, 1992.
62. Purity M, Hever-Szabo A and Venetianer A: Overexpression of P-glycoprotein in heat- and/or drug-resistant hepatoma variants. *Cytotechnology* 19:207-214, 1996.
63. Pratt WB: The role of the hsp90-based chaperone system in signal transduction by nuclear receptors and receptors signaling via MAP kinase. *Annu Rev Pharmacol Toxicol* 37:297-326, 1997.
64. Pratt WB and Toft DO: Steroid receptor interactions with heat shock protein and immunophilin complexes. *Endocrine Rev* 18:306-360, 1997.
65. Punyiczki M and Fésüs L: Heat shock and apoptosis: the two defense systems of the organisms may have overlapping molecular elements. *Ann NY Acad Sci* 851:67-74, 1998.
66. Santarosa M, Favaro D, Quaià M, et al: Expression of heat shock protein 72 in renal cell carcinoma: possible role and prognostic implications in cancer patients. *Eur J Cancer* 33:873-877, 1997.
67. Santin AD, Hermonat PL, Ravaggi A, et al: The effects of irradiation on the expression of a tumour rejection antigen (heat shock protein gp96) in human cervical cancer. *Int J Radiat Biol* 73:699-704, 1998.
68. Schirmer EC, Glover JR, Singer MA, et al: HSP100/Clp proteins: a common mechanism explains diverse functions. *Trends Biochem Sci* 21:289-296, 1996.
69. Schweighoffer T: Tumor cells expressing a recall antigen are powerful cancer vaccines. *Eur J Immunol* 26:2559-2564, 1996.
70. Seo SJ, Park YM, Kim JI, et al: T-cell lymphoma in transgenic mice expressing hsp 70 gene. *Biochem Biophys Res Commun* 218:582-587, 1996.
71. Song HY, Dunbar JD, Zhang YX, et al: Identification of a protein with homology to hsp90 that binds the type 1 tumor necrosis factor receptor. *J Biol Chem* 270:3574-3581, 1995.
72. Spector NL, Hardy L, Ryan C, et al: 28-kDa mammalian heat shock protein, a novel substrate of a growth regulatory protease involved in differentiation of human leukemia cells. *J Biol Chem* 270:1003-1006, 1995.
73. Srivastava PK: Endo- β -D-glucuronidase (heparanase) activity of heat-shock protein/tumour rejection antigen gp96. *Biochem J* 301:918, 1994.
74. Srivastava PK and Heike M: Tumor-specific immunogenicity of stress-induced proteins: convergence of two evolutionary pathways of antigen presentation? *Semin Immunol* 3:57-64, 1991.
75. Srivastava PK and Maki RG: Stress-induced proteins in immune response to cancer. *Curr Top Microbiol Immunol* 167:109-123, 1991.
76. Srivastava PK, DeLeo AB and Old LJ: Tumor rejection antigens of chemically induced tumors of inbred mice. *Proc Natl Acad Sci USA* 83:3407-3411, 1986.
77. Srivastava PK, Udono H, Blachere NE, et al: Heat shock proteins transfer peptides during antigen processing and CTL priming. *Immunogenetics* 39:93-98, 1994.
78. Stahl J, Wobus AM, Ithrig S, et al: The small heat shock protein hsp25 is accumulated in P19 embryonal carcinoma cells and embryonic stem cells of line BLC6 during differentiation. *Differentiation* 51:33-37, 1992.
79. Stepanova L, Leng X, Parker SB, et al: Mammalian p50Cdc37 is a protein kinase-targeting subunit of Hsp90 that binds and stabilizes Cdk4. *Genes Dev* 10:1491-1502, 1996.
80. Storm FK, Mahvi DM and Gilchrist KW: Heat shock protein 27 overexpression in breast cancer lymph node metastasis. *Ann Surg Oncol* 3:570-573, 1996.
81. Sugawara S, Takeda K, Lee A, et al: Suppression of stress protein GRP78 induction in tumor B/C10ME eliminates resistance to cell mediated cytotoxicity. *Cancer Res* 53:6001-6005, 1993.
82. Sullivan DM, Ben-Yosef R and Kapp DS: Stanford 3D hyperthermia treatment planning system. Technical review and clinical summary. *Int J Hyperthermia* 9:627-643, 1993.
83. Suto R and Srivastava PK: A mechanism for the specific immunogenicity of heat shock protein-chaperoned peptides. *Science* 269:1585-1588, 1995.
84. Tamura Y, Peng P, Liu K, et al: Immunotherapy of tumors with autologous tumor-derived heat shock protein preparations. *Science* 278:117-120, 1997.
85. Udono H and Srivastava PK: Heat shock protein-peptide complexes, reconstituted in vitro, elicit peptide-specific cytotoxic T lymphocyte response and tumor immunity. *J Exp Med* 186:1315-1322, 1997.
86. Udono H, Levey DL and Srivastava PK: Cellular requirements for tumor-specific immunity elicited by heat shock proteins: tumor rejection antigen gp96 primes CD8+ T cells in vivo. *Proc Natl Acad Sci USA* 91:3077-3081, 1994.
87. Ullrich SJ, Robinson EA, Lav LW, et al: A mouse tumor-specific transplantation antigen is a heat-shock related protein. *Proc Natl Acad Sci USA* 83:3121-3125, 1986.
88. van Eden W and Young DB (eds): *Stress Proteins in Medicine*. Marcel Dekker Inc, New York, 1996.
89. Walker RA, Jones JL, Chappell S, et al: Molecular pathology of breast cancer and its application to clinical management. *Cancer Metastasis Rev* 16:6-27, 1997.
90. Wang G, Klostergaard J, Khodadadian M, et al: Murine cells transfected with human Hsp27 cDNA resist TNF-induced cytotoxicity. *J Immunother Emphasis Tumor Immunol* 19:9-20, 1996.
91. Wawrzynow A, Banecki B and Zyllicz M: The Clp ATPases define a novel class of molecular chaperones. *Mol Microbiol* 21:895-899, 1996.
92. Welch WJ: Mammalian stress response: cell physiology, structure/function of stress proteins, and implications for medicine and disease. *Physiol Rev* 72:1063-1081, 1992.
93. Yano M, Naito Z, Tanaka S, et al: Expression and roles of heat shock proteins in human breast cancer. *Jpn J Cancer Res* 87:908-915, 1996.
94. Yufu Y, Nishimura J and Nawata H: High constitutive expression of heat shock protein 90 α in human acute leukemia cells. *Leuk Res* 16:597-605, 1992.